

PERSISTENT CURRENTS IN A KONDO RING

H.-P. Ecker,^{1,2} H. Johannesson,³ and C. A. Stafford^{2,4}

¹*Department of Physics, University of Jyväskylä, FIN-40351 Jyväskylä, Finland*

²*Department of Physics, University of Arizona, 1118 E. 4th Street, Tucson, AZ 85721, USA*

³*Institute of Theoretical Physics, Chalmers University of Technology and Göteborg University, SE 412 96 Göteborg, Sweden*

⁴*Fakultät für Physik, Albert-Ludwigs-Universität, Hermann-Herder-Straße 3, D-79104 Freiburg, Germany*

The influence of a magnetic impurity or ultrasmall quantum dot on the persistent current of a mesoscopic ring is investigated. The system consists of electrons in a one-dimensional ring threaded by an Aharonov-Bohm flux Φ , coupled via an antiferromagnetic exchange interaction to a localized electron. The problem is mapped onto a Kondo model for the even-parity channel plus free electrons in the odd-parity channel. The twisted boundary conditions couple states of opposite parity unless $\Phi = f\Phi_0/2$, where $\Phi_0 = hc/e$ is the flux quantum and f is an integer. For these special values of Φ , the model is solved by the Bethe ansatz, and it is shown that the charge stiffness is insensitive to the presence of the magnetic impurity/quantum dot.

1 Introduction

Mesoscopic Kondo physics is a subject of considerable current interest, both experimentally^{1,2,3} and theoretically.^{1,4} It has recently become possible to measure Kondo scattering from a single magnetic impurity.² And in an experimental tour de force, electron transport through a quantum dot in the Kondo regime has also been observed.³ With Kondo physics in quantum dots now accessible in the laboratory, the full mesoscopic richness of this paradigm of many-body physics can be investigated. One of the cornerstones of mesoscopia is the Aharonov-Bohm (AB) effect, which has already been observed in microstructured rings coupled to a quantum dot.⁵ It is natural to ask how the AB effect in a ring coupled to a quantum dot would be modified by the many-body correlations present in the Kondo regime.

The equilibrium response of a multiply-connected system to an AB flux Φ piercing it is the persistent current.^{6,7,8,9} The persistent current of a ring coupled via tunneling to a quantum dot was investigated via perturbation theory and numerical diagonalization by Büttiker and Stafford.⁹ In this article, we study a variant of the problem where electrons in a one-dimensional (1D) ring threaded by an AB flux are coupled via antiferromagnetic exchange to a localized electron, representing a magnetic impurity or quantum dot. A detailed analysis shows that this model can be mapped onto the integrable Kondo model for special values of Φ , corresponding to periodic and antiperiodic boundary conditions. The model is solved via Bethe ansatz for these special values of Φ , and it is shown that the charge stiffness is insensitive to the Kondo scattering, showing that spin-charge separation holds even on the mesoscopic scale in this model.

2 Exactly solvable model

The system we are contemplating is described in the continuum limit by the 1D Hamiltonian,

$$H = -\frac{\hbar^2}{2m} \sum_{\alpha} \int_0^L dx \psi_{\alpha}^{\dagger}(x) \partial_x^2 \psi_{\alpha}(x) + \lambda \sum_{\alpha, \beta} \psi_{\alpha}^{\dagger}(0) \vec{\sigma}_{\alpha\beta} \psi_{\beta}(0) \cdot \vec{S}, \quad (1)$$

where $\lambda > 0$ is an antiferromagnetic Kondo coupling, m is the electron mass, L the ring's circumference, \vec{S} is the impurity spin (located at $x = 0$), and ψ_{α} is an electron field with spin index $\alpha = \uparrow, \downarrow$. The effect of the magnetic flux Φ has been gauged away⁶ and encapsulated in twisted boundary conditions:

$$\psi_{\alpha}(L) = e^{i\phi} \psi_{\alpha}(0), \quad (2)$$

where $\phi = 2\pi\Phi/\Phi_0$, with $\Phi_0 = hc/e$ the elementary flux quantum.

We are interested in an exact solution of the problem described by (1) and (2), with a particular eye on how the Kondo interaction in (1) may affect the persistent current induced by the flux Φ . Since the essential physics of the system is confined to a small region around the left and right Fermi points, we can linearize the quadratic dispersion in (1) around $\pm k_F$ and introduce left (l) and right (r) moving chiral fields:

$$\psi_\alpha(x) \sim e^{-ik_F x} \psi_{l,\alpha}(x) + e^{ik_F x} \psi_{r,\alpha}(x). \quad (3)$$

The Hamiltonian then becomes:

$$H = H_0 + H_{\text{imp}}, \quad (4)$$

with

$$H_0 = \frac{v_F}{2\pi} \sum_\alpha \int_0^L dx \left(\psi_{l,\alpha}^\dagger(x) i \partial_x \psi_{l,\alpha}(x) - \psi_{r,\alpha}^\dagger(x) i \partial_x \psi_{r,\alpha}(x) \right), \quad (5)$$

and

$$H_{\text{imp}} = \lambda \sum_{\alpha,\beta} \left(\psi_{l,\alpha}^\dagger(0) + \psi_{r,\alpha}^\dagger(0) \right) \vec{\sigma}_{\alpha\beta} \left(\psi_{l,\beta}(0) + \psi_{r,\beta}(0) \right) \cdot \vec{S}. \quad (6)$$

To make progress, it is convenient to pass to a basis of definite parity fields (*Weyl basis*):

$$\psi_{\text{even},\alpha}(x) = \frac{1}{\sqrt{2}} (\psi_{r,\alpha}(x) + \psi_{l,\alpha}(-x)), \quad (7)$$

an even-parity, right-moving electron field, and

$$\psi_{\text{odd},\alpha}(x) = \frac{1}{\sqrt{2}} (\psi_{r,\alpha}(-x) - \psi_{l,\alpha}(x)), \quad (8)$$

an odd-parity, left-moving field. One should note that the assignment of chirality (*left/right*) to parity (*odd/even*) is not intrinsic, but a property of the particular transformations (7) and (8). This is analogous to a gauge-fixing condition. In this basis, the Hamiltonian takes the form:

$$H = H_0^{\text{odd}} + H_0^{\text{even}} + H_{\text{imp}}^{\text{even}}, \quad (9)$$

where

$$H_0^{\text{even}} = -\frac{v_F}{2\pi} \sum_\alpha \int_0^L dx \psi_{\text{even},\alpha}^\dagger(x) i \partial_x \psi_{\text{even},\alpha}(x) \quad (10)$$

and

$$H_0^{\text{odd}} = \frac{v_F}{2\pi} \sum_\alpha \int_0^L dx \psi_{\text{odd},\alpha}^\dagger(x) i \partial_x \psi_{\text{odd},\alpha}(x) \quad (11)$$

describe independent relativistic electrons, and the impurity contribution is now also diagonal:

$$H_{\text{imp}}^{\text{even}} = \lambda \sum_{\alpha,\beta} \psi_{\text{even},\alpha}^\dagger(0) \vec{\sigma}_{\alpha\beta} \psi_{\text{even},\beta}(0) \cdot \vec{S}. \quad (12)$$

We recognize $H_K^{\text{even}} \equiv H_0^{\text{even}} + H_{\text{imp}}^{\text{even}}$ as the chiral Hamiltonian of the spin- S Kondo model.

While the even and odd parity channels are decoupled in the Hamiltonian, there is a price to be paid: the twisted boundary conditions (2) couple states of opposite parity:

$$\begin{pmatrix} \psi_{\text{even},\alpha}(L) \\ \psi_{\text{odd},\alpha}(L) \end{pmatrix} = \begin{pmatrix} \cos \phi & i \sin \phi \\ -i \sin \phi & \cos \phi \end{pmatrix} \begin{pmatrix} \psi_{\text{even},\alpha}(0) \\ \psi_{\text{odd},\alpha}(0) \end{pmatrix}, \quad (13)$$

where in (3) we have taken $k_F = (2\pi/L)n$, with n an integer. For the special values $\phi = f\pi$ (i.e., $\Phi = f\Phi_0/2$), where f is an integer, the matrix in Eq. (13) reduces to a multiple of the unit

matrix, and the even and odd parity states decouple from each other entirely. One can then solve H_K^{even} by the Bethe ansatz.¹⁰ Thus, our original problem in (1) and (2) has collapsed to an exactly solvable problem for $f \in Z$, consisting of a left-moving odd-parity branch of independent relativistic electrons, together with a (decoupled) right-moving even-parity branch defined by the 1D Kondo model. For generic values of ϕ , it is not possible to choose a basis which renders the Hamiltonian and the boundary conditions simultaneously diagonal, strongly suggesting that the model is not integrable in general. This is in apparent contradiction to recent claims in the literature¹³ about the integrability of the related Anderson ring threaded by an Aharonov-Bohm flux of arbitrary strength.

From Eq. (12), the impurity is seen to couple only to the spin current of the electrons, suggesting, via the dynamic spin-charge separation in 1D, that the (charge) persistent current is insensitive to the presence of the impurity. Although this indeed turns out to be the case—as we shall confirm via a Bethe ansatz analysis—some caveats are appropriate at this point: First, the persistent current is a boundary effect and, as such, could be influenced by *non-dynamical* selection rules for combining charge and spin.^{11,12} Secondly, and possibly reflecting this, a magnetic impurity *does* affect the charge current of a *chiral* ring of free electrons (with all electrons moving in the same direction).¹⁴ In any event, it is instructive to study the exact mechanism by which the charge persistent current in the present problem avoids any influence from the impurity.

To carry out this analysis, we first need to consider how to properly define a persistent current for relativistic electrons, i.e. for electrons with a *linear* dispersion.

3 Persistent Current for Relativistic Electrons

In the usual treatment of independent 1D electrons,^{7,8} the persistent current is obtained by summing the partial currents $I_n = -(e/\hbar)\partial E_n/\partial\phi$ over all occupied levels n . This approach clearly fails for the relativistic electrons in (10) and (11), since the corresponding linear dispersions

$$E_r = \hbar v_F \frac{2\pi n_r + \phi}{L}, \quad n_r = 0, 1, 2, \dots, n_F \quad (14)$$

and

$$E_l = \hbar v_F \frac{-2\pi n_l + \phi}{L}, \quad n_l = 1, 2, \dots, n_F \quad (15)$$

imply that $\partial E_n/\partial\phi = \text{const.}$ for all levels n . (Here we consider a system of spinless electrons in which the total number of electrons $2n_F + 1$ is odd.) To recover the known results for the persistent current, we must thus use an alternative approach. Let us introduce *flux-dependent* particle numbers

$$N_{r/l}(\phi) = \frac{L}{2\pi} [|k_{r/l,F}(\phi)| - |k_{r/l,F}(0)|], \quad (16)$$

where $k_{r/l,F}$ are flux-dependent Fermi momenta, connected to the highest occupied level on the respective branch. $k_{r,F}$ and $k_{l,F}$ are cutoff dependent, and need not be equal, since only the right-movers couple to the impurity in Eq. (9). However, provided the cutoffs are chosen independent of ϕ , $N_{r/l}(\phi)$ are insensitive to the cutoff, and describe the physical response of the system to an AB flux. The persistent current is then

$$I(\phi) = -\frac{ev_F}{L} [N_r(\phi) - N_l(\phi)]. \quad (17)$$

It should be pointed out that the charge velocity v_F is in general subject to renormalization due to interactions.

With the choice of representative levels in (14) and (15) (note in particular that the zero mode of the original problem (1) is assigned to *one* branch only) it is easy to verify that (16)

and (17) exactly reproduce the known result for the persistent current of an odd number of spinless 1D electrons.⁸ Our construction, introduced here *ad hoc*, can be put on a firm basis by a proper analysis of the cutoff procedure for 1D relativistic electrons in the presence of an Aharonov-Bohm flux.¹⁵ In short, a flux-dependent particle number as in (16) is the trade-off that guarantees that physical observables remain independent of the choice of cutoff (bounding the spectrum from below).

Given (16) and (17), the problem is now reduced to calculating how the Fermi momenta $k_{r/l,F}$ depend on the flux and the coupling of the electrons to the magnetic impurity. For this, we turn to a finite-size Bethe ansatz analysis.

4 Finite-size Bethe ansatz

To obtain the Fermi momenta $k_{r/l,F}(\phi)$ for a finite ring, we apply the techniques of the *Bethe ansatz* for finite systems, developed previously for the 1D Hubbard model.¹⁶ As pointed out above, our model is only integrable for $\phi = f\pi$, with f an integer. For $f \in \mathbb{Z}$, the nested Bethe ansatz equations which diagonalize H are

$$Lk_{n_l} = -2\pi n_l + f\pi + \frac{2\pi}{N_{\text{odd}}} \sum_{\delta=1}^{M_{\text{odd}}} J_{\delta}, \quad (18)$$

$$Lk_{n_r} = 2\pi n_r + f\pi + \sum_{\gamma=1}^{M_{\text{even}}} [\Theta(2\Lambda_{\gamma} - 2) - \pi], \quad (19)$$

$$N_{\text{even}}\Theta(2\Lambda_{\gamma} - 2) + \Theta(2\Lambda_{\gamma}) = 2\pi I_{\gamma} + \sum_{\delta=1}^{M_{\text{even}}} \Theta(\Lambda_{\gamma} - \Lambda_{\delta}), \quad (20)$$

where k_{n_l} are the pseudomomenta characterizing the N_{odd} odd-parity left movers which decouple from the impurity, M_{odd} of which have spin down, and k_{n_r} are pseudomomenta characterizing the N_{even} even-parity right movers, M_{even} of which have spin down, and $\{\Lambda_{\gamma}, \gamma = 1, \dots, M_{\text{even}}\}$ are a set of auxiliary variables known as spin-rapidities. The scattering phase shifts are given by $\Theta(x) = -\tan^{-1}(x/c)$, with $c = 2\lambda/(1 - 3\lambda/4)$. Eq. (18) simply gives the quantum numbers of free, chiral electrons, written in the Bethe basis. The Bethe ansatz equation (19) describes the charge degrees of freedom in the even channel (holons), while Eq. (20) describes the spin degrees of freedom in the even channel (spinons). Eqs. (19) and (20) differ from the Bethe ansatz equations derived previously for the Kondo model¹⁰ only by the addition of the AB flux $\phi = f\pi$.

The persistent current is an odd function of ϕ by symmetry,⁶ and is analytic, except at values of ϕ corresponding to level crossings. We are interested in the persistent current for small values of the AB flux. Choosing the total numbers of both up- and down-spin electrons to be odd excludes a level crossing at $\phi = 0$. The leading mesoscopic behavior of the persistent current is then

$$I(\phi) = -D_c\phi/L + \mathcal{O}(\phi^3/L^3), \quad (21)$$

where D_c is the charge stiffness. Eq. (21) holds on general grounds independent of whether the model is integrable or not.

The choice of quantum numbers $\{n_l, J_{\delta}, n_r, I_{\gamma}\}$ specifies the quantum state of the system. Generically, there are one or more level crossings¹⁷ between $f = 0$ and $f = 1$. To determine the charge stiffness, however, we only need to consider the state which evolves adiabatically from the ground state at $f = 0$ as ϕ is increased. This state is given by $M_{\text{even/odd}} = (N_{\text{even/odd}} + / - 1)/2$, (with $N_{\text{even/odd}}$ odd for simplicity), with integer-spaced quantum numbers $\{n_l, J_{\delta}, n_r, I_{\gamma}\}$ in the symmetric ranges $-(N_{\text{odd}} - 1)/2 \leq n_l \leq (N_{\text{odd}} - 1)/2$, $-(M_{\text{odd}} - 1)/2 \leq J_{\delta} \leq (M_{\text{odd}} - 1)/2$, $-(N_{\text{even}} - 1)/2 \leq n_r \leq (N_{\text{even}} - 1)/2$, and $-(M_{\text{even}} - 1)/2 \leq I_{\gamma} \leq (M_{\text{even}} - 1)/2$. The quantum

numbers of the even-parity sector are the same as those of the Kondo model with periodic boundary conditions.¹⁰

Given a set of spin rapidities Λ_γ satisfying Eq. (20), we may calculate the sum in Eq. (19), and thus the momenta k_{n_r} are determined. One sees immediately that the total scattering phase shift of the dressed magnetic impurity is *independent* of f , so that $N_r = -N_l = f/2$. In addition, the charge velocity v_F is unrenormalized by interactions in this model.¹⁰ The charge stiffness may be evaluated from Eqs. (17) and (21) as a finite difference $D_c = -LI(f=1)/\pi = ev_F/\pi + \mathcal{O}(L^{-2})$, and is unaffected by the Kondo scattering. The persistent current for small Φ is thus

$$I = -\frac{ev_F}{L} \frac{2\Phi}{\Phi_0}, \quad (22)$$

which is identical to the result for free electrons. Eq. (22) indicates that spin-charge separation holds even at the mesoscopic scale in this model.

Acknowledgments

This work was supported in part by a bilateral grant from the Finnish Academy and the Deutscher Akademischer Austauschdienst. HPE is an Alexander-von-Humboldt Fellow of the Finnish Academy. He thanks the Department of Physics of the University of Arizona and the Institute of Theoretical Physics at Chalmers and Göteborg University for kind hospitality during crucial stages of this work. HJ acknowledges financial support from the Swedish Natural Science Research Council.

References

1. For a review, see *Physics Today*, January 1998, page 17, and references therein.
2. J. Li, W.-D. Schneider, R. Berndt, and R. Delley, *Phys. Rev. Lett.* **80**, 2893 (1998).
3. D. Goldhaber-Gordon's contribution to these proceedings.
4. See, e.g., L. Glazman's contribution to these proceedings.
5. A. Yacoby, M. Heiblum, D. Mahalu, and H. Shtrikman, *Phys. Rev. Lett.* **74**, 4047 (1995).
6. N. Byers and C. N. Yang, *Phys. Rev. Lett.* **7**, 46 (1961).
7. M. Büttiker, Y. Imry, and R. Landauer, *Phys. Lett.* **A96**, 365 (1983).
8. H.-F. Cheung *et al.*, *Phys. Rev. B* **37**, 6050 (1988).
9. M. Büttiker and C. A. Stafford, *Phys. Rev. Lett.* **76**, 495 (1996).
10. For a recent review, see: N. Andrei, *Integrable Models in Condensed Matter Physics*, in *Series on Modern Condensed Matter Physics - Vol. 6*, 458-551 (World Scientific, Singapore, 1992), Eds. S. Lundquist, G. Morandi, and Yu Lu.
11. P. Fröjdh and H. Johannesson, *Phys. Rev. B* **53**, 3211 (1996).
12. C. A. Stafford, *Phys. Rev. B* **48**, 8430 (1993).
13. P. Schlottmann and A. A. Zvyagin, *Phys. Lett. A* **231**, 109 (1997).
14. A. A. Zvyagin and T. V. Bandos, *Low Temp. Phys.* **20**, 222 (1994).
15. H.-P. Eckle, H. Johannesson and C. A. Stafford, (unpublished).
16. F. Woynarovich and H.-P. Eckle, *J. Phys. A: Math. Gen.* **20**, L443 (1987); F. Woynarovich, *J. Phys. A: Math. Gen.* **22**, 2615 (1989).
17. N. Yu and M. Fowler, *Phys. Rev. B* **45**, 11795 (1992); C. A. Stafford and A. J. Millis, *Phys. Rev. B* **48**, 1409 (1993).